

| Ordnungen und Gattungen der fossilen Flora von Bilin. | Zahl der neuen Arten. | Gesamtzahl der Arten. | Vorkommen in anderen fossi l. Floren. | | | | |
|---|--------------------------|--------------------------|---------------------------------------|---------|---|--------|--------|
| | | | Tertiärflora der Schweiz | Radoboj | Niederrheini- sche Braun- kohlenflora | Häring | Sotzka |
| Ord: Sapotaceae. | | 10 | 8 | 3 | 2 | 9 | 5 |
| <i>Sapotacites</i> | 2 | 5 | 7 | — | — | 8 | 4 |
| <i>Chrysophyllum</i> | 2 | 2 | — | — | 1 | — | — |
| <i>Bumelia</i> | 2 | 3 | 1 | 2 | 1 | 1 | 1 |
| „ Ebenaceae. | | 7 | 3 | 7 | 1 | 1 | — |
| <i>Diospyros</i> | 3 | 5 | 2 | 5 | 1 | 1 | — |
| <i>Macreightia</i> | 1 | 2 | 1 | — | — | — | — |
| „ Styraceae. | | 2 | 1 | 3 | — | — | — |
| <i>Styrax</i> | 1 | 2 | 1 | 2 | — | — | — |
| „ Vaccinieae. | | 1 | 10 | 2 | 1 | — | 2 |
| <i>Vaccinium</i> | | 1 | 10 | 2 | 1 | — | 2 |
| „ Ericaceae. | | 7 | 9 | 5 | 2 | 3 | 1 |
| <i>Arbutites</i> | 1 | 1 | — | — | — | — | — |
| <i>Andromeda</i> | 2 | 3 | 4 | 2 | 1 | 2 | 1 |
| <i>Azalea</i> | 1 | 2 | — | 1 | — | — | — |
| <i>Rhododendron</i> | 1 | 1 | — | 1 | — | — | — |